

NATIONAL BUREAU OF STANDARDS REPORT

1602

SOME RELATIONS AMONG THE BLOCKS OF SYMMETRICAL
GROUP DIVISIBLE DESIGNS

by

W. S. CONNOR



U. S. DEPARTMENT OF COMMERCE
NATIONAL BUREAU OF STANDARDS



THE NATIONAL BUREAU OF STANDARDS

The scope of activities of the National Bureau of Standards is suggested in the following listing of the divisions and sections engaged in technical work. In general, each section is engaged in specialized research, development, and engineering in the field indicated by its title. A brief description of the activities, and of the resultant reports and publications, appears on the inside of the back cover of this report.

1. **ELECTRICITY.** Resistance Measurements. Inductance and Capacitance. Electrical Instruments. Magnetic Measurements. Electrochemistry.
2. **OPTICS AND METROLOGY.** Photometry and Colorimetry. Optical Instruments. Photographic Technology. Length. Gage.
3. **HEAT AND POWER.** Temperature Measurements. Thermodynamics. Cryogenics. Engines and Lubrication. Engine Fuels.
4. **ATOMIC AND RADIATION PHYSICS.** Spectroscopy. Radiometry. Mass Spectrometry. Physical Electronics. Electron Physics. Atomic Physics. Neutron Measurements. Nuclear Physics. Radioactivity. X-Rays. Betatron. Nucleonic Instrumentation. Radiological Equipment. Atomic Energy Commission Instruments Branch.
5. **CHEMISTRY.** Organic Coatings. Surface Chemistry. Organic Chemistry. Analytical Chemistry. Inorganic Chemistry. Electrodeposition. Gas Chemistry. Physical Chemistry. Thermochemistry. Spectrochemistry. Pure Substances.
6. **MECHANICS.** Sound. Mechanical Instruments. Aerodynamics. Engineering Mechanics. Hydraulics. Mass. Capacity, Density, and Fluid Meters.
7. **ORGANIC AND FIBROUS MATERIALS.** Rubber. Textiles. Paper. Leather. Testing and Specifications. Organic Plastics. Dental Research.
8. **METALLURGY.** Thermal Metallurgy. Chemical Metallurgy. Mechanical Metallurgy. Corrosion.
9. **MINERAL PRODUCTS.** Porcelain and Pottery. Glass. Refractories. Enameled Metals. Building Stone. Concreting Materials. Constitution and Microstructure. Chemistry of Mineral Products.
10. **BUILDING TECHNOLOGY.** Structural Engineering. Fire Protection. Heating and Air Conditioning. Exterior and Interior Coverings. Codes and Specifications.
11. **APPLIED MATHEMATICS.** Numerical Analysis. Computation. Statistical Engineering. Machine Development.
12. **ELECTRONICS.** Engineering Electronics. Electron Tubes. Electronic Computers. Electronic Instrumentation.
13. **ORDNANCE DEVELOPMENT.** Mechanical Research and Development. Electromechanical Fuzes. Technical Services. Missile Fuzing Research. Missile Fuzing Development. Projectile Fuzes. Ordnance Components. Ordnance Tests. Ordnance Research.
14. **RADIO PROPAGATION.** Upper Atmosphere Research. Ionospheric Research. Regular Propagation Services. Frequency Utilization Research. Tropospheric Propagation Research. High Frequency Standards. Microwave Standards.
15. **MISSILE DEVELOPMENT.** Missile Engineering. Missile Dynamics. Missile Intelligence. Missile Instrumentation. Technical Services. Combustion.

NBS PROJECT

NBS REPORT

1163-11-1107

17 April 1952

1662

**SOME RELATIONS AMONG THE BLOCKS OF SYMMETRICAL
GROUP DIVISIBLE DESIGNS**

by

W. S. Connor



This report is issued
in any form, either in
from the Office of the

Approved for public release by the
Director of the National Institute of
Standards and Technology (NIST)
on October 9, 2015

tion, reprinting, or reproduction
permission in writing is obtained
Washington 25, D. C.

THE REPORT

1953-1954

1953-1954

1953-1954

1953-1954

1953-1954

1953-1954

1953-1954

1953-1954

1953-1954

1953-1954

1953-1954

1953-1954

1953-1954

1953-1954

1953-1954

FOREWORD

The combinatorial problem which is studied in this paper is of interest in the design of experiments.

The paper considers the way in which the blocks of a symmetrical Group Divisible incomplete block design, with parameters v, r, m, n, λ_1 , and λ_2 , are connected by common treatments. Structural and characteristic matrices are defined, and a relation among them is exhibited. It is shown that if a solution exists for a given set of parameters which are such that $r \neq \lambda_1$, $r^2 \neq v\lambda_2$, and $r^2 - v\lambda_2$ and $\lambda_1 - \lambda_2$ are relatively prime, then the interchange of blocks with treatments yields a solution which corresponds to the given set of parameters.

J. H. Curtiss
Chief, National Applied
Mathematics Laboratories

A. V. Astin
Acting Director
National Bureau of Standards

FOREWORD

This paper is of interest in the design of experiments. The combinatorial problem which is studied is

The paper considers the way in which the blocks of a symmetric group divisible incomplete block design with parameters v, r, k, λ and λ_1 are connected by common treatments. Structural and characteristic matrices are defined, and a relation among them is exhibited. It is shown that if a solution exists for a given set of parameters which are such that v, k, λ, λ_1 and r are relatively prime, then the interchange of blocks with treatments yields a solution which corresponds to the given set of parameters.

J. H. GILLIS
Chief, National Applied
Mathematics Laboratories

A. V. HALL
Acting Director
National Bureau of Standards

SOME RELATIONS AMONG THE BLOCKS OF SYMMETRICAL GROUP DIVISIBLE DESIGNS

W. S. Connor¹
National Bureau of Standards

1. SUMMARY. It is well known that if every pair of treatments in a symmetrical Balanced incomplete block design occurs in λ blocks, then every two blocks of the design have λ treatments in common. In this paper it will be shown that a somewhat similar property holds for symmetrical Group Divisible designs. In the course of the investigation there will be introduced certain matrices which are of intrinsic interest.

2. INTRODUCTION. Some of the combinatorial properties of Group Divisible incomplete block designs were considered in [1]. Here we shall need the definition of Group Divisible designs and the three classes into which they fall. An incomplete block design with v treatments each replicated r times in b blocks of size k is said to be Group Divisible (GD) if the treatments can be divided into m groups, each with n treatments, so that the treatments belonging to the same group occur together in λ_1 blocks and the treatments belonging to different groups occur together in λ_2 blocks, $\lambda_1 \neq \lambda_2$. The three exhaustive and mutually exclusive classes into which the GD designs fall are as follows:

- (a) Singular GD designs characterized by $r - \lambda_1 = 0$;
- (b) Semi-regular GD designs characterized by $r - \lambda_1 > 0$.

¹This work was begun while the author was at the University of North Carolina.

SOME RELATIONS AMONG THE BLOCKS OF SYMMETRIC GROUP DIVISIBLE DESIGNS

W. S. CONNOR
National Bureau of Standards

1. SUMMARY. It is well known that if every pair of

treatments in a symmetrical balanced incomplete block design occurs in λ blocks, then every two blocks of the design have a treatment in common. In this paper it will be shown that a somewhat similar property holds for symmetrical Group Divisible

designs. In the course of the investigation there will be introduced certain matrices which are of intrinsic interest.

2. INTRODUCTION. Most of the combinatorial properties

of Group Divisible incomplete block designs were considered in [1]. Here we shall now consider the definition of Group Divisible

designs and the three classes into which they fall: an incomplete block design with v treatments each replicated r times in b blocks of size k is said to be Group Divisible (GD)

if the treatments can be divided into m groups, each of size s , such that the k treatments belonging to the same group occur together in λ blocks and the treatments belonging to different groups occur together in λ' blocks. λ, λ', s, k are integers. The three exhaustive and mutually exclusive classes into which the GD designs fall are as follows:

- (a) Singular GD designs characterized by $r = \lambda = 0$.
- (b) Semi-regular GD designs characterized by $r = \lambda$.

$rk - v\lambda_2 = 0$; and

(c) Regular GD designs characterized by $r - \lambda_1 > 0$,
 $rk - v\lambda_2 > 0$.

In this paper we shall study classes (b) and (c) for the symmetrical case, that is, the case when $r = k$, or equivalently, $b = v$.

3. THE INCIDENCE AND STRUCTURAL MATRICES. In [2] there was defined the structural matrix for Balanced incomplete block designs. We now shall define the incidence matrix, and two structural matrices for GD designs.

Let us consider first the incidence matrix of a GD design,

$$(3.1) \quad N = \begin{bmatrix} n_{11} & \cdots & n_{1b} \\ \vdots & & \vdots \\ n_{v1} & \cdots & n_{vb} \end{bmatrix},$$

where the rows represent treatments, the columns represent blocks, and $n_{ij} = 1$ or 0 according as the i -th treatment does or does not occur in the j -th block. From the conditions satisfied by the design it is easy to see that

$$(3.2) \quad \sum_{j=1}^b n_{ij} = r, \quad (i = 1, \dots, v),$$

and

$$(3.3) \quad \sum_{j=1}^b n_{ij}n_{uj} = \lambda_1 \text{ or } \lambda_2,$$

$$x - \sqrt{2} = 0; \text{ and}$$

(c) Regular GD designs characterized by $x - \sqrt{2} = 0$.

$$x - \sqrt{2} > 0.$$

In this paper we shall study classes (b) and (c) for

the symmetrical case, that is, the case when $x = 1$, or

$$\text{equivalently, } v = 2.$$

2. THE INCIDENCE AND STRUCTURAL MATRICES. In [2] there

was defined the structural matrix for balanced incomplete block

designs. We now shall define the incidence matrix, and the

structural matrices for GD designs.

Let us consider first the incidence matrix of a GD design.

$$(2.1) \quad N = \begin{bmatrix} n_{11} & n_{12} & \dots & n_{1p} \\ n_{21} & n_{22} & \dots & n_{2p} \\ \vdots & \vdots & \ddots & \vdots \\ n_{v1} & n_{v2} & \dots & n_{vp} \end{bmatrix}$$

where the rows represent treatments, the columns represent

blocks, and $n_{ij} = 1$ or 0 according as the i -th treatment does

or does not occur in the j -th block. From the conditions

satisfied by the design it is easy to see that

$$(2.2) \quad \sum_{j=1}^p n_{ij} = r, \quad (i = 1, \dots, v).$$

and

$$(2.3) \quad \sum_{i=1}^v n_{ij} = k, \quad (j = 1, \dots, p).$$

according as the i -th and u -th treatments ($i \neq u$) do belong or do not belong to the same group.

Throughout the paper let us adopt the convention that the treatments $n(w-1) + 1, n(w-1) + 2, \dots, nw$ shall belong to the w -th group, ($w=1, \dots, n$). Then

$$(3.4) \quad NN' = \begin{bmatrix} A & B & \circ & \circ & \circ & B \\ B & A & \circ & \circ & \circ & B \\ \circ & \circ & & & & \circ \\ \circ & \circ & & & & \circ \\ \circ & \circ & & & & \circ \\ \circ & \circ & & & & \circ \\ B & B & \circ & \circ & \circ & A \end{bmatrix},$$

where the elements of the $n \times n$ submatrix A are r in the principal diagonal and λ_1 elsewhere, and the elements of the $n \times n$ submatrix B are λ_2 everywhere. Of course NN' contains $v = mn$ rows and columns.

Now choose any $t \leq b$ blocks of the design. Let the submatrix of N which corresponds to these t blocks be denoted by N_0 . Let s be the number of treatments common to the j -th and u -th chosen blocks, ($j, u = 1, 2, \dots, t$). Then the $t \times t$ symmetric matrix

$$(3.5) \quad S_t^I = N_0' N_0 = (s_{ju})$$

is defined to be the intersection structural matrix of the t chosen blocks. The j -th row or column of S_t corresponds to the



j -th chosen block and the successive elements of the j -th row or column give the number of treatments which this block has in common with the 1st, 2nd, ..., t -th chosen blocks.

We next shall consider another structural matrix. Let s_{ju}^w denote the number of treatments from the w -th group which blocks j and u have in common. Then

$$(3.6) \quad \sum_{w=1}^m s_{ju}^w = s_{ju}.$$

and

$$(3.7) \quad \sum_{w=1}^m s_{jj}^w = k.$$

Now consider the matrix

$$(3.8) \quad G_t = \begin{bmatrix} 1 & 1 & \dots & 1 \\ s_{11}^1 & s_{22}^1 & \dots & s_{tt}^1 \\ 2 & 2 & \dots & 2 \\ s_{11}^2 & s_{22}^2 & \dots & s_{tt}^2 \\ \vdots & \vdots & \ddots & \vdots \\ m & m & \dots & m \\ s_{11}^m & s_{22}^m & \dots & s_{tt}^m \end{bmatrix},$$

and the product matrix

$$(3.9) \quad s_t^G = G_t^t G_t.$$

where the element in the j -th row and u -th column is the sum of

6. 1. 16. 1978

... ..
... ..
... ..
... ..
... ..
... ..

... ..
... ..
... ..

... ..
... ..
... ..

... ..
... ..
... ..

... ..

... ..

... ..

products of the number of treatments which the j -th chosen block and the u -th chosen block contain from each group. We define S_t^G as the group structural matrix of the t chosen blocks.

4. THE CHARACTERISTIC MATRIX. We shall define an analogue of the characteristic matrix which was developed for Balanced incomplete block designs in [2]. For the remainder of the paper, except for the last section, we shall restrict our attention to the regular GD designs.

Let the columns of N be permuted so that the first t columns correspond to the t chosen blocks. Then let the incidence matrix be extended by adjoining t new rows, so that the elements of the j -th adjoined row are zero, except for the j -th which is unity. We thus get

$$(4.1) \quad N_1 = \begin{bmatrix} N \\ I_t & 0 \end{bmatrix},$$

where ~~which~~ I_t is the identity matrix of order t , and 0 is the $t \times (b-t)$ zero matrix. Then

$$(4.2) \quad N_1 N_1^0 = \begin{bmatrix} N N^0 & N 0 \\ N^0 0 & I_t \end{bmatrix}$$

The evaluation of $|N_1 N_1^0|$ leads to

The first part of the document discusses the importance of maintaining accurate records of all transactions. It emphasizes that every entry, no matter how small, should be recorded to ensure the integrity of the financial data. The second part of the document provides a detailed breakdown of the various types of transactions that may occur, including sales, purchases, and transfers. It also outlines the specific steps that should be followed to record each type of transaction accurately. The third part of the document discusses the importance of reconciling the records on a regular basis to identify any discrepancies and correct them as soon as possible. The fourth part of the document provides a summary of the key points discussed and offers some final thoughts on the importance of maintaining accurate records.

Summary of Key Points	
1. Maintain accurate records of all transactions.	2. Record every transaction, no matter how small.
3. Breakdown transactions into sales, purchases, and transfers.	4. Follow specific steps to record each type of transaction.
5. Reconcile records regularly to identify discrepancies.	6. Correct discrepancies as soon as possible.
7. Summarize key points and offer final thoughts.	8. Emphasize the importance of maintaining accurate records.

$$(4.3) \quad |N_1 N_1^t| = (rk)^{-t+1} (r - \lambda_1)^{v-t-m} (rk - v\lambda_2)^{m-t-1} |C_t| ,$$

where the typical element of C_t is

$$(4.4) \quad c_{ju} = (rk - v\lambda_2) (rk \delta_{ju} + \lambda_2 k^2) \\ + (\lambda_1 - \lambda_2) (rk \sum_{w=1}^m s_{jj}^w s_{uu}^w - n \lambda_2 k^2) ,$$

where $\delta_{ju} = (r - \lambda_1 - k)$ or $-\delta_{ju}$, according as $j = u$ or $j \neq u$.

The matrix C_t is defined as the characteristic matrix of the t chosen blocks. The j -th row or the j -th column of C_t corresponds to the j -th chosen block of the design.

We observe that the characteristic matrix is related to the two structural matrices as is described in the following theorem.

Theorem 4.1. For the regular GD designs there exists a (1,1) correspondence among the elements of the intersection structural matrix S_t^I , the group structural matrix S_t^G , and the characteristic matrix C_t . This correspondence is given by

$$C_t = rk(rk - v\lambda_2) [(r - \lambda_1) I_t - S_t^I] + rk(\lambda_1 - \lambda_2) S_t^G \\ + \lambda_2 k^2 (r - \lambda_1) E_t ,$$

where I_t is the identity matrix of order t , E_t is the singular

$t \times t$ matrix all of whose elements are unity, and the other quantities are scalars.

For the particular case when $r = k$, the value of $|N_1 N_1|$ as given by (4.3) reduces to

$$(4.5) \quad |N_1 N_1| = r^{-2}(t-1)(r-\lambda_1)^{v-t-m}(r^2-v\lambda_1)^{n-t-1}|C_t|,$$

where the typical element of C_t is

$$(4.6) \quad c_{ju} = r^2(r^2-v\lambda_2)(\delta_{ju}+\lambda_2) + r^2(\lambda_1-\lambda_2)\left(\sum_{j=1}^m s_j s_{ju}^{v-n\lambda_2}\right).$$

We shall state an analogue of Theorem 3.1 of [2]. The proof is as for that theorem.


Theorem 4.2. If C_t is the characteristic matrix of any set of t blocks chosen from a regular GD design with parameters $v, b, r, k, m, n, \lambda_1$, and λ_2 , then

- (i) $|C_t| \geq 0$ if $t < b-v$,
- (ii) $|C_t| = 0$ if $t > b-v$, and
- (iii) $r^{-2}(t-1)(r-\lambda_1)^{2v-b-m}(r^2-v\lambda_2)^{n-b+v-1}|C_{b-v}|$

is a perfect integral square.

5. INEQUALITIES ON s_{ju} FOR REGULAR SYMMETRICAL GD DESIGNS.

Let $t=1$. Then since the factor outside of $|C_1|$ in (4.5) is positive, it follows from Theorem 4.2 that $|C_1| = 0$. Hence from



(4.6),

$$(5.1) \quad r^2(\lambda_1 - \lambda_2) \left[\sum_{w=1}^m (s_{11}^w)^2 - r^2 - v\lambda_2 - n\lambda_2 \right] = 0.$$

Since $r^2(\lambda_1 - \lambda_2) \neq 0$,

$$(5.2) \quad \sum_{w=1}^m (s_{11}^w)^2 = r^2 - v\lambda_2 + n\lambda_2$$

Now let $t = 2$. Since $c_{11} = c_{22} = 0$, it is necessary by Theorem 4.2 that $c_{12} = c_{21} = 0$. Hence from (4.6)

$$(5.3) \quad s_{12} = \lambda_2 + \frac{e}{(r^2 - v\lambda_2)} (\lambda_1 - \lambda_2),$$

where

$$e = \sum_{w=1}^m s_{11}^w s_{22}^w - n\lambda_2.$$

From (5.2) and the observation that $s_{jj}^w \geq 0$, ($j=1,2$; $w=1, \dots, m$), it follows that

$$(5.4) \quad -n\lambda_2 \leq e \leq r^2 - v\lambda_2.$$

From (5.3) and (5.4) we obtain

Theorem 5.1. For a regular symmetrical CD design the number of treatments s_{ju} common to two blocks satisfies the inequalities

$$\lambda_2(r - \lambda_1) / (r^2 - v\lambda_2) \leq s_{ju} \leq \lambda_1,$$

when $\lambda_1 > \lambda_2$. The inequalities are reversed when $\lambda_1 < \lambda_2$.



6. THE BLOCK STRUCTURE FOR REGULAR SYMMETRICAL GD DESIGNS
WHEN $r^2 - v\lambda_2$ AND $\lambda_1 - \lambda_2$ ARE RELATIVELY PRIME. We need to consider the distribution of the treatments contained in an initial block B_1 among the other blocks. Let n_j be the number of blocks among the remaining $(b-1)$ blocks which has j treatments in common with B_1 . Then from the definition of the design we obtain

$$(6.1) \quad \sum_{j=0}^k n_j = b-1 = v-1, \text{ and}$$

$$\sum_{j=0}^k j n_j = r(k-1) = r(r-1).$$

Also consider $M = \sum_{j=0}^k j(j-1)n_j$, which is twice the number of pairs of treatments of B_1 which lie among the other blocks.

M is given by

$$(6.2) \quad M = \sum_{w=1}^n s_{11}^w (s_{11}^w - 1) (\lambda_1 - 1) + \sum_{\substack{x, w=1 \\ x \neq w}}^n s_{11}^x s_{11}^w (\lambda_2 - 1).$$

From (3.7) and (5.2), since $r=k$,

$$(6.3) \quad \sum_{w=1}^n s_{11}^w (s_{11}^w - 1) = (n-1) \lambda_1, \text{ and}$$

$$(6.4) \quad \sum_{\substack{x, w=1 \\ x \neq w}}^n s_{11}^x s_{11}^w = (n-1) n \lambda_2.$$

Hence

$$(6.5) \quad M = (n-1) (\lambda_1) (\lambda_1 - 1) + (n-1) (n) (\lambda_2) (\lambda_2 - 1).$$

The first part of the paper is devoted to the study of the properties of the function $f(x)$ defined by the equation $f(x) = \sum_{n=0}^{\infty} \frac{x^n}{n!}$. It is shown that $f(x)$ is a continuous function and that it satisfies the differential equation $f'(x) = f(x)$. The second part of the paper is devoted to the study of the properties of the function $g(x)$ defined by the equation $g(x) = \sum_{n=0}^{\infty} \frac{x^n}{n!} \ln n$. It is shown that $g(x)$ is a continuous function and that it satisfies the differential equation $g'(x) = g(x) + \frac{1}{x}$.

The third part of the paper is devoted to the study of the properties of the function $h(x)$ defined by the equation $h(x) = \sum_{n=0}^{\infty} \frac{x^n}{n!} \ln^2 n$. It is shown that $h(x)$ is a continuous function and that it satisfies the differential equation $h'(x) = h(x) + \frac{2}{x}$. The fourth part of the paper is devoted to the study of the properties of the function $k(x)$ defined by the equation $k(x) = \sum_{n=0}^{\infty} \frac{x^n}{n!} \ln^3 n$. It is shown that $k(x)$ is a continuous function and that it satisfies the differential equation $k'(x) = k(x) + \frac{3}{x}$.

The fifth part of the paper is devoted to the study of the properties of the function $l(x)$ defined by the equation $l(x) = \sum_{n=0}^{\infty} \frac{x^n}{n!} \ln^4 n$. It is shown that $l(x)$ is a continuous function and that it satisfies the differential equation $l'(x) = l(x) + \frac{4}{x}$. The sixth part of the paper is devoted to the study of the properties of the function $m(x)$ defined by the equation $m(x) = \sum_{n=0}^{\infty} \frac{x^n}{n!} \ln^5 n$. It is shown that $m(x)$ is a continuous function and that it satisfies the differential equation $m'(x) = m(x) + \frac{5}{x}$.

Now consider

$$(6.6) \quad B = \sum_{j=0}^k (j - \lambda_1)(j - \lambda_2) n_j .$$

From (6.1), (6.5), and (6.6) we obtain

$$(6.7) \quad B = 0 .$$

Hence the following lemma.

Lemma 6.1. If for a regular symmetrical GD design n_j denotes the number of blocks which have j treatments in common with a given initial block, then

$$B = \sum_{j=0}^k n_j (j - \lambda_1)(j - \lambda_2) = 0 .$$

Now let $r^2 - v\lambda_2$ and $\lambda_1 - \lambda_2$ be relatively prime. It follows from (5.3) that s_{12} cannot lie in the open interval (λ_1, λ_2) . Then every term of B is positive or zero. But since $B = 0$, every term must be zero. We thus get

Theorem 6.1. If for a regular symmetrical GD design $r^2 - v\lambda_2$ and $\lambda_1 - \lambda_2$ are relatively prime, then any two blocks have either λ_1 or λ_2 treatments in common.

We further observe that even if $r^2 - v\lambda_2$ and $\lambda_1 - \lambda_2$ are not relatively prime, it still may not be possible to choose the elements of G_t of (3.8), subject to the restrictions of (3.7) and (5.2), such that s_{ju} is integral, *but is not λ_1 or λ_2 .* Consider, for

example, the GD design with parameters $v=b=15$, $r=k=9$, $m=3$, $n=15$, $\lambda_1=3$, and $\lambda_2=1$. The H.C.F. of $r^2-v\lambda_2$ and $\lambda_1-\lambda_2$ is 2. It is clear that the only positive integers which satisfy (3.7) and (5.2) are 1, 1, and 7. But then we must have either $\sum_{j=1}^m s_{jj}^v s_{uu}^v = 51$ or 15, which correspond respectively to λ_1 and λ_2 .

Now assume that the condition of theorem 6.1 is met, or more generally, that positive integers do not exist which meet the restrictions of (3.7), (5.2) and Lemma 6.1 and imply values of s_{ju} other than λ_1 and λ_2 . Then from (6.1), we obtain

$$\begin{aligned} (6.8) \quad n\lambda_1 + n\lambda_2 &= v-1, \text{ and} \\ \lambda_1 n\lambda_1 + \lambda_2 n\lambda_2 &= r(r-1), \end{aligned}$$

whence

$$\begin{aligned} (6.9) \quad n\lambda_1 &= n-1, \text{ and} \\ n\lambda_2 &= (n-1)n, \end{aligned}$$

so that with respect to any initial block B_1 , there are $(n-1)$ other blocks which have λ_1 treatments in common with it, and $(n-1)n$ other blocks which have λ_2 treatments in common with it.

From (5.3) we see that

$$(6.10) \quad \sum_{j=1}^m s_{11}^v s_{jj}^v = r + (n-1)\lambda_1$$

implies that blocks i and j have λ_1 treatments in common, and

conversely. But then from (5.2) and (6.10), it follows that

$$(6.11) \quad \sum_{w=1}^n s_{11}^w s_{jj}^w = \sum_{w=1}^n (s_{11}^w)^2,$$

which implies that $s_{11}^w = s_{jj}^w$, ($w=1, \dots, n$; $j=2, \dots, b$). Hence, if blocks B_1 and B_j have λ_1 treatments in common, and blocks B_1 and B_u have λ_1 treatments in common, then B_j and B_u have λ_1 treatments in common. We thus have

Theorem 6.2. If for a regular symmetrical GD design $r^2 - v\lambda_2$ and $\lambda_1 - \lambda_2$ are relatively prime, then the blocks fall into n groups of n blocks each, which are such that any two blocks from the same group contain λ_1 treatments in common and any two blocks from different groups contain λ_2 treatments in common.

As has been indicated above, this theorem could be stated somewhat more generally.

7. THE SEMI-REGULAR CLASS. For this class ~~it was shown~~ ^{$rk - v\lambda_2 = 0$, and} in [1] that ~~$s_{jj}^w = k$, a constant, for all w and j .~~ Hence the above theory does not apply. We shall give a simple example which demonstrates for small v that there do sometimes exist solutions in which $s_{ju} \neq \lambda_1$ or λ_2 for some j and u .

Consider the GD design with parameters $v=b=8$, $r=k=4$, $n=4$, $n=2$, $\lambda_1 = 0$, and $\lambda_2 = 2$. One solution is

$$N^{(1)} = \begin{bmatrix} 1 & 1 & 1 & 1 & 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 & 1 & 1 & 1 & 1 \\ 1 & 1 & 0 & 0 & 0 & 0 & 1 & 1 \\ 0 & 0 & 1 & 1 & 1 & 1 & 0 & 0 \\ 0 & 1 & 1 & 0 & 1 & 0 & 1 & 0 \\ 1 & 0 & 0 & 1 & 0 & 1 & 0 & 1 \\ 1 & 0 & 1 & 0 & 0 & 1 & 1 & 0 \\ 0 & 1 & 0 & 1 & 1 & 0 & 0 & 1 \end{bmatrix},$$

which has the property, that the blocks break up into 4 groups of 2 blocks each, such that two blocks in the same group have zero treatments in common and any two blocks from different groups have 2 treatments in common.

Another solution is

$$N^{(2)} = \begin{bmatrix} 1 & 1 & 1 & 1 & 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 & 1 & 1 & 1 & 1 \\ 1 & 0 & 0 & 1 & 1 & 0 & 0 & 1 \\ 0 & 1 & 1 & 0 & 0 & 1 & 1 & 0 \\ 0 & 1 & 0 & 1 & 0 & 1 & 0 & 1 \\ 1 & 0 & 1 & 0 & 1 & 0 & 1 & 0 \\ 0 & 0 & 1 & 1 & 0 & 0 & 1 & 1 \\ 1 & 1 & 0 & 0 & 1 & 1 & 0 & 0 \end{bmatrix},$$

which is such that any initial block has 1 treatment in common with each of three blocks, 2 treatments in common with each of three blocks, and 3 treatments in common with one block.

We now shall obtain inequalities for the number of treatments λ_{10} common to any two blocks of a symmetrical semi-regular GD design. Since for a semi-regular GD design, $rk = v\lambda_2$, it follows that $r - \lambda_1 = n(\lambda_2 - \lambda_1)$, from which we obtain the following lemma.

Theorem 4.1. For a semi-regular GD design, it is necessary that $\lambda_2 \geq \lambda_1$.

Now let $r = k$. Choose any two blocks and let the columns of N be permuted so that the first two columns correspond to the chosen blocks. Then to N affix a new column, the v -th of which contains $(\lambda_2 - \lambda_1)^{1/2}$ in the rows which correspond to the treatments of the two blocks, $(1, 2, \dots, n)$, and zero elsewhere. Let the augmented matrix be denoted by N_2 . Now form

$$(7.1) \quad N_2 = \begin{bmatrix} N_2 & \\ I_2 & 0 \end{bmatrix}$$

where I_2 is the identity matrix of order 2 and 0 is the $2 \times (v-2)$ matrix all of whose elements are zero. Then

$$(7.2) \quad |N_2 N_2'| = (r+v\lambda_2 - \lambda_1)^{-1} (r - \lambda_1)^{v-3} |B_2|$$

where B_2 is a 2×2 matrix with elements

$$(7.3) \quad \begin{aligned} b_{11} &= b_{22} = (r+v\lambda_2 - \lambda_1)(-\lambda_1) + \lambda_2 r^2, \text{ and} \\ b_{12} &= b_{21} = (r+v\lambda_2 - \lambda_1)(-\lambda_{12}) + \lambda_1 r^2. \end{aligned}$$

As for Theorem 4.2 it is necessary that $|N_2 N_2'| \geq 0$, and since the factor outside of $|B_2|$ in (7.2) is positive, it is necessary that $|B_2| \geq 0$. Hence, the following theorem:

Theorem 7.1. For a symmetrical semi-regular GD design, the number of treatments common to two blocks, s_{ju} , satisfies the inequalities

$$1 \leq s_{ju} \leq \frac{2\lambda_1 r^2}{r-v\lambda_2-\lambda_1} - \lambda_1$$

I wish to express my thanks ^{to} ~~for~~ Professor R. C. Bose for suggesting this problem.

REFERENCES

- [1] R. C. Bose and W. S. Connor, "Combinatorial Properties of Group Divisible Incomplete Block Designs", Annals of Mathematical Statistics, submitted for publication.
- [2] W. S. Connor, Jr., "On the Structure of Balanced Incomplete Block Designs", Annals of Mathematical Statistics, Vol. 23, (1952), pp. 57-71.

Reports and Publications

The results of the Bureau's research for the development of equipment and devices for publication are published in various forms. Published reports appear either in the Bureau's own journals or in other publications or in the journals of other organizations. The Bureau itself publishes technical reports, which are available from the Government Printing Office. The Bureau also publishes technical reports, which are available from the Government Printing Office. The Bureau also publishes technical reports, which are available from the Government Printing Office. The Bureau also publishes technical reports, which are available from the Government Printing Office.

Information on the Bureau's publications can be obtained from the Bureau's publications section. Information on the Bureau's publications can be obtained from the Bureau's publications section. Information on the Bureau's publications can be obtained from the Bureau's publications section. Information on the Bureau's publications can be obtained from the Bureau's publications section.

EXHIBIT

11. H. T. Jones and W. H. Jones, "The Influence of the Group on the Individual's Moral Behavior," Psychological Studies, Vol. 1, No. 1, 1932, pp. 1-10.
12. H. T. Jones and W. H. Jones, "The Influence of the Group on the Individual's Moral Behavior," Psychological Studies, Vol. 1, No. 1, 1932, pp. 1-10.

THE NATIONAL BUREAU OF STANDARDS

Functions and Activities

The National Bureau of Standards is the principal agency of the Federal Government for fundamental and applied research in physics, mathematics, chemistry, and engineering. Its activities range from the determination of physical constants and properties of materials, the development and maintenance of the national standards of measurement in the physical sciences, and the development of methods and instruments of measurement, to the development of special devices for the military and civilian agencies of the Government. The work includes basic and applied research, development, engineering, instrumentation, testing, evaluation, calibration services, and various scientific and technical advisory services. A major portion of the NBS work is performed for other government agencies, particularly the Department of Defense and the Atomic Energy Commission. The functions of the National Bureau of Standards are set forth in the Act of Congress, March 3, 1901, as amended by Congress in Public Law 619, 1950. The scope of activities is suggested in the listing of divisions and sections on the inside of the front cover.

Reports and Publications

The results of the Bureau's work take the form of either actual equipment and devices or published papers and reports. Reports are issued to the sponsoring agency of a particular project or program. Published papers appear either in the Bureau's own series of publications or in the journals of professional and scientific societies. The Bureau itself publishes three monthly periodicals, available from the Government Printing Office: the Journal of Research, which presents complete papers reporting technical investigations; the Technical News Bulletin, which presents summary and preliminary reports on work in progress; and Basic Radio Propagation Predictions, which provides data for determining the best frequencies to use for radio communications throughout the world. There are also five series of nonperiodical publications: the Applied Mathematics Series, Circulars, Handbooks, Building Materials and Structures Reports, and Miscellaneous Publications.

Information on the Bureau's publications can be found in NBS Circular 460, Publications of the National Bureau of Standards (\$1.00). Information on calibration services and fees can be found in NBS Circular 483, Testing by the National Bureau of Standards (25 cents). Both are available from the Government Printing Office. Inquiries regarding the Bureau's reports and publications should be addressed to the Office of Scientific Publications, National Bureau of Standards, Washington 25, D. C.

